








# Addressing Stress and Rhythm Dysregulation among Health Workers in the Post-Pandemic Context: An Italian Study and a Proposal for a Preventive Psychoeducational Intervention



Elisa Cantone<sup>1,2,\*</sup> , Antonio Urban<sup>1</sup> , Cesar Aviles Gonzales<sup>3,4</sup> , Alessandra Perra<sup>2,\*</sup> , Stefano Lorrai<sup>2</sup>, Michela Atzeni<sup>2</sup> , Stocchino Serena<sup>2</sup>, Rober Romero Ramirez<sup>3</sup>, Adalberto Rangel Restrepo<sup>3</sup>, Mauro Giovanni Carta<sup>2</sup>  and Giulia Cossu<sup>2</sup> 

<sup>1</sup>Capacities Building for Global Health, University of Cagliari, 09124 Cagliari, Italy

<sup>2</sup>Department of Medical Sciences and Public Health, University of Cagliari, 09124 Cagliari, Italy

<sup>3</sup>Alumni Group of Nursing, Universidad Popular del Cesar, Valledupar 200002, Colombia

<sup>4</sup>Department of Medicine and Surgery, University of Enna "Kore", Piazza dell'Università, 94100 Enna, Italy

## Abstract:

**Introduction:** During the COVID-19 pandemic, healthcare workers (HCWs) faced intense and prolonged stress. This exposure increased their risk of burnout and mood disorders. The *Dysregulation of Mood, Energy, and Social Rhythms (DYMERS) hypothesis* suggests that instability in biological and social rhythms may precede the onset of mood dysregulation. This study examined rhythm disturbances among HCWs and explored the potential role of a rhythm-focused psychoeducational intervention.

**Methods:** Ninety-seven HCWs from the University Hospital of Cagliari, Italy, participated in a cross-sectional study. Depressive symptoms were assessed using the Patient Health Questionnaire-9 (PHQ-9), and rhythm regulation was evaluated through the Biological Rhythms Interview of Assessment in Neuropsychiatry (BRIAN). Data were compared with published findings from mood disorders, psychotic disorders, and community samples.

**Results:** A current depressive episode with PHQ-9 score was identified in 35.1% of HCWs participants, with significantly higher prevalence ( $p = 0.01$ ) among women (42.7%) than men (14.7%). Mean BRIAN scores were  $46.1 \pm 11.0$ , significantly higher than those observed in psychotic disorder ( $p = 0.04$ ) and community samples ( $p < 0.0001$ ). Rhythm dysregulation scores approached values reported in bipolar disorder samples. Elevated BRIAN scores were also observed among non-depressed participants.

**Discussion:** The elevated rhythm dysregulation scores observed in healthcare workers support the hypothesis that chronic occupational stress may disrupt biological and social rhythms, potentially increasing vulnerability to mood disorders.

**Conclusion:** HCWs displayed pronounced rhythm dysregulation and high rates of depressive symptoms, highlighting occupational stress as a potential major risk factor and further supporting the DYMERS hypothesis. From both clinical and research perspectives, it could be essential to promote psychoeducational programs that target rhythm regulation, such as adaptations of Interpersonal and Social Rhythm Therapy, which may offer effective preventive strategies to enhance resilience and mental well-being in this population. However, the cross-sectional design, convenience sampling, limited socio-demographic data, and reliance on self-reported measures constrain generalizability and causal inference, underscoring the need for larger longitudinal studies to validate and extend these findings.

**Keywords:** Healthcare workers, DYMERS, COVID-19, Rhythm dysregulation, Prevention, Depressive symptoms.

© 2026 The Author(s). Published by Bentham Open.

This is an open access article distributed under the terms of the Creative Commons Attribution 4.0 International Public License (CC-BY 4.0), a copy of which is available at: <https://creativecommons.org/licenses/by/4.0/legalcode>. This license permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.



Received: September 18, 2025  
Revised: November 13, 2025  
Accepted: November 19, 2025  
Published: February 27, 2026



Send Orders for Reprints to  
[reprints@benthamscience.net](mailto:reprints@benthamscience.net)

\*Address correspondence to these authors at the Capacities Building for Global Health, University of Cagliari, 09124 Cagliari, Italy and Department of Medical Sciences and Public Health, University of Cagliari, 09124 Cagliari, Italy;  
E-mails: [elisa.cantone@libero.it](mailto:elisa.cantone@libero.it) and [alessandra.perra@unica.it](mailto:alessandra.perra@unica.it)

Cite as: Cantone E, Urban A, Gonzales C, Perra A, Lorrà S, Atzeni M, Serena S, Ramírez R, Restrepo A, Carta M, Cossu G. Addressing Stress and Rhythm Dysregulation among Health Workers in the Post-Pandemic Context: An Italian Study and a Proposal for a Preventive Psychoeducational Intervention. *Clin Pract Epidemiol Ment Health*, 2026; 22: e17450179449051. <http://dx.doi.org/10.2174/0117450179449051260114063307>

## 1. INTRODUCTION

The *Social Zeitgeber Hypothesis* (from the German *zeitgeber*, “time giver”) proposes that disruptions in social routines can destabilize biological rhythms [1]. These routines include sleep-wake cycles, mealtimes, and interpersonal interactions. In vulnerable individuals, such disturbances may trigger mood episodes [2, 3].

Evidence shows that life events disrupting daily routines and sleep instability often precede depressive or (hypo)manic episodes [4, 5]. Similarly, sleep instability and circadian disruption are commonly observed in individuals with Bipolar Disorder (BD) even during euthymic phases [6-8]. Importantly, therapeutic approaches designed to stabilize rhythm, such as Interpersonal and Social Rhythm Therapy (IPSRT), have demonstrated efficacy in improving remission and psychosocial functioning [9, 10]. Collectively, these findings emphasize rhythm regulation as a critical mechanism in both the onset and management of BD. Building on this framework, the concept of DYMERS (“Dysregulation of Mood, Energy, and Social Rhythms Syndrome”) was recently introduced [11]. A preclinical condition is characterized by behavioral rhythm dysregulation that may increase vulnerability to mood disorders depending on individual stress exposure. DYMERS has been linked as a transdiagnostic vulnerability factor. It could affect clinical populations with different disorders such as mood disorders, panic disorder, attention deficit/hyperactivity disorder (ADHD), and post-traumatic stress disorder (PTSD) [12-14]. However, it could also affect high-risk groups, including healthcare workers (HCWs) and individuals with chronic illnesses, who may experience burnout and other stress-related outcomes [15-17]. This underscores the broader clinical relevance of rhythm dysregulation in vulnerable populations.

HCWs are particularly vulnerable to rhythm dysregulation due to night shifts, unpredictable schedules, understaffing, and organizational strain. Evidence shows that misalignment between internal circadian rhythms and work schedules increases the risk of burnout, depression, and poor sleep, independent of shift type [18-22]. The COVID-19 pandemic further exacerbated occupational stress [23], making HCWs an ideal population to investigate the presence and impact of rhythm disruption in a non-clinical, high-risk context.

### 1.1. Study Aim

Against this background, the present study aimed to investigate mood and rhythm dysregulation in HCWs, a population at high risk for mental health difficulties due to post-pandemic stress, workload, and organizational instability

The specific objectives were to:

1. Assess the prevalence and severity of depressive symptoms among HCWs using a standardized self-report instrument.
2. Evaluate the degree of biological rhythm dysregulation in the same sample through a validated international measure.
3. Examine the relationship between depressive symptoms and rhythm dysregulation, exploring whether greater rhythm instability is associated with current depressive episodes.
4. Compare rhythm dysregulation levels in HCWs with those previously reported in clinical samples with bipolar and psychotic disorders, and in a community sample from the same geographic area.
5. Discuss the implications of these findings for the development of rhythm-focused psychoeducational interventions aimed at prevention and early support in this high-risk occupational group.

In this non-clinical but high-risk population, we hypothesize that HCWs will exhibit biological and social rhythm dysregulation at levels comparable to clinical populations with mood disorders. The co-occurrence of depressive symptoms among those with rhythm dysregulation may indicate increased vulnerability for future mood episodes, consistent with the Social Zeitgeber and DYMERS frameworks. Given the cross-sectional design, no causal conclusions can be drawn, but identifying both rhythm dysregulation and depressive symptoms provides important heuristic insight for future longitudinal and preventive studies.

## 2. METHODS

### 2.1. Design

This study employed a cross-sectional design to investigate rhythm dysregulation and depressive symptoms among HCWs. Participants were recruited from four outpatient facilities affiliated with the University Hospital of

Cagliari, Italy, providing a representative sample of health professionals working in diverse clinical settings. Voluntary convenience sampling will be conducted based on the availability of eligible participants at each recruiting center, following verification of inclusion criteria and acquisition of informed consent. The study was conducted and reported in accordance with the STROBE statement guidelines for observational studies [24].

## 2.2. Study Sample

Participants eligible for inclusion in the study were adults aged 18 years and older, of any gender, who provided signed informed consent. Eligible healthcare workers were those professionally engaged in care services within specified categories, including physicians, nurses, psychologists, healthcare assistants, social workers, and rehabilitation staff. Individuals who did not meet these inclusion criteria or who had severe medical or psychiatric conditions that would prevent them from completing the study questionnaires were excluded. Data collection occurred during a single week in May 2025 to capture a consistent snapshot of participants' status. The sample included 97 HCWs, comprising both fully qualified staff and trainees who, according to Italian regulations, are authorized to perform clinical duties during their final postgraduate years. This approach ensured that all participants had active responsibilities in healthcare delivery, making the findings relevant to real-world clinical practice. Given the voluntary nature of participation, a potential self-selection bias cannot be excluded, as individuals experiencing higher levels of stress or workload may have been less likely to participate.

## 2.3. Study Instruments

Following participants' signing of informed consent, which was approved by the local ethics committee, the following validated instruments were administered:

### 2.3.1. Demographic and Work-related Questionnaire

This tool collected essential background information, such as age, gender, employment ward, and professional role, to characterize the sample and explore potential associations with study outcomes. Some socio-demographic data, such as years of service and type of work shifts, could represent potential sources of bias, since participant recruitment was conducted within a single healthcare organization. For this reason, collecting such data might have risked compromising participant anonymity and violating privacy principles.

### 2.3.2. Patient Health Questionnaire - 9 item (PHQ-9)

The Italian version of this widely used self-report scale [25, 26] was utilized to screen for depressive episodes. The PHQ-9 aligns with DSM-5 diagnostic criteria by assessing nine core depressive symptoms. Scores range from 0 to 27, with a cut-off of 10 or above indicating a high probability of clinically relevant depressive symptoms, typically reflecting mild to moderate depression severity. This tool's psychometric properties and clinical utility are well established internationally.

### 2.3.3. Biological Rhythms Interview of Assessment in Neuropsychiatry (BRIAN)

To evaluate disruptions in biological and social rhythms, we used the Italian validated version of BRIAN [27, 28]. This semi-structured interview assesses five key rhythm domains: sleep patterns, physical activity, social interactions, meal timing, and chronotype (a measure of an individual's preferred timing of daily activities). The questionnaire consists of 21 items rated on a four-point scale (1 = "not at all" to 4 = "often"), with higher scores indicating greater rhythm dysregulation. BRIAN has demonstrated strong reliability and has been translated and applied across multiple cultural contexts [29, 30].

## 2.4. Statistical Analysis

All data analyses were performed with IBM SPSS Statistics software (version 28.0.1, IBM Corp., Armonk, NY, USA). Descriptive statistics were calculated to summarize participant characteristics and study variables. Categorical variables were analyzed using Chi-square tests with Yates' correction when necessary, and non-parametric methods were applied when distributional assumptions were not met. Normality of continuous variables (BRIAN scores) was verified through the Shapiro-Wilk test. Given approximate normality, parametric analyses were performed using one-way ANOVA with Bonferroni post-hoc tests, reporting Cohen's  $d$  as the effect size for pairwise comparisons. To account for demographic differences, BRIAN scores were adjusted for age and sex prior to comparison with published reference samples. Differences between the present sample and previously published samples were then statistically tested using independent-sample t-tests.

## 3. RESULTS

A total of 97 HCWs completed the study assessments and were included in the sample. Two invited workers (one nurse and one medical doctor) refused participation (2.1% of the overall sample), citing pressing work commitments that prevented them from completing the study instruments. The sample represents a group of HCWs experiencing significant psychological distress (Table 1). Specifically, thirty-four participants (35.05%) met criteria for a current depressive episode. No significant differences emerged between participants above and below 50 years of age. Women showed a higher prevalence of depressive symptoms compared to men, reaching statistical significance ( $\chi^2 = 5.76$ ,  $p = 0.016$ ), while non-physicians exhibited a non-significant trend toward higher depression rates than physicians (48.9% vs. 22.9%,  $p = 0.059$ ). The mean BRIAN score for the sample was  $46.14 \pm 10.96$ , indicating moderate biological rhythm dysregulation. Women had higher mean scores than men ( $47.64 \pm 12.82$  vs.  $42.62 \pm 9.68$ ), but the difference did not reach statistical significance ( $p = 0.059$ ). Participants with current depressive episodes reported markedly higher BRIAN scores compared to participants without current depressive episodes ( $55.12 \pm 7.94$  vs.  $41.29 \pm 12.58$ ;  $p < 0.001$ ;  $d = 1.3$ ).

**Table 1. Characteristics of the sample, frequency of depressive episodes, and BRIAN score.**

	HCWS (N=97)	PHQ<10 N=63 (64.95%)	PHQ>9 N=34 (35.05%)	STAT
<b>Men and Female N(%)</b>	29 (29.9%) 68 (70.1%)	24 (85.3%) 39 (57.35%)	5 (14.7%) 29 (42.65%)	<b>Depressive Episode by sex:</b> $\chi^2 = 5.764$ $p=0.016$ or women = 3.57 ci95%1.2-10.5
<b>&gt;49 and &lt;50 N(%)</b>	36 (37.11%) 61 (62.89%)	23 (63.89%) 40 (65.57%)	13 (36.11%) 21 (34.43%)	<b>Depressive Episode by age:</b> $\chi^2 = 0.013$ $p=0.10$ or <50 = 0.95 ci95%0.4-2.2
<b>Mean Age</b>	43.89±12.64	43.74±12.95	44.17±11.50	1,95df $f=0.026$ , $p=0.872$
<b>Medical Doctors and Not Medical N(%)</b>	35 (36.08%) 62 (63.92%)	27 (77.14%) 36 (51.06)	8 (22.86%) 26 (48.94%)	<b>Depressive Episode by role:</b> $\chi^2 = 3.577$ $p=0.059$ or notmd = 2.44 ci95%1.0-6.2
<b>Mean Brian Score</b>	46.14±10.96	41.29±12.58	55.12±7.94	<b>Difference between depressed and not depressed:</b> 1,95df $f=37.740$ $p<0.0001$
<b>Men and Female Mean Brian Score</b>	42.62±9.68 47.64±12.82	40.01±10.88 41.83±13.30	52.60±3.88 55.55±8.64	<b>Difference by Sex:</b> 1,95df $f=3.569$ $p=0.062$

**Table 2. Comparison with a published study on the BRIAN Score in clinical and non-clinical samples in the same area.**

N(%)	Stabilized Psychosis [31] (N=51)	BD [32] (N=40)	Community Sample [27] (N=82)	BD [33] (N=64)
<b>Men</b>	40 (65.57%)	12 (30%)	32 (39.0%)	21 (32.8%)
<b>Mean Age</b>	37.22±10.7	47.23 ± 13.37	44.63±12.95	47.23 ± 13.37
<b>Brian Score</b>	40.40±7.02	45.07 ± 12.00	22.22±11.19	48.97±12.54
<b>Brian Score</b>	43.70±10.36	46.29±9.76	44.77±11.59	45.99±10.71
<b>HCWS Standardized</b>				
<b>ANOVA 1 Way</b>	1,146 df	1,135 df	1,177 df	1,159 df
	$f=4.162$	$f=0.386$	$f=173.602$	$f=2.603$
	$p=0.043$	$p=0.536$	$p<0.0001$	$p=0.109$
	$d=0.37$		$d=1.97$	

When compared with previously published data from the same geographic area (Table 2), the mean BRIAN score observed among HCWs did not significantly differ from that of individuals with bipolar disorder reported in two independent studies ( $p = 0.536$ ;  $p = 0.109$ ). Conversely, it was significantly higher than that observed in a sample of individuals with stabilized psychosis ( $43.70 \pm 10.36$  vs.  $40.40 \pm 7.02$ ;  $p = 0.043$ ;  $d = 0.37$ ) and markedly higher than in a community sample ( $44.77 \pm 11.59$  vs.  $22.22 \pm 11.19$ ;  $p < 0.001$ ;  $d = 1.97$ ).

**4. DISCUSSION**

The findings of this study reveal a concerning degree of biological rhythm dysregulation among HCWs, a population that remains understudied within the context of rhythm-related psychopathology [34, 35]. Notably, the mean BRIAN scores observed in this non-clinical yet highly stressed group closely resemble those reported in individuals

diagnosed with BD in previous research. Although the mean rhythm dysregulation observed among HCWs approached the levels reported in clinical populations, this should not be interpreted as diagnostic equivalence. Instead, these findings suggest a subclinical rhythm vulnerability consistent with the DYMERS framework, rather than a fully prodromal condition [36, 37]. The high prevalence of current depressive episodes (35.05%), particularly among women, aligns with global trends in post-pandemic mental health and underscores the central role of rhythm instability in mood deterioration [38-40]. These results support the conceptualization of rhythm disruptions not merely as secondary manifestations of mood disorders, but as core transdiagnostic mechanisms that may precede the onset of formal psychiatric diagnoses [13].

Furthermore, BRIAN scores in our sample exceeded those reported for individuals with psychosis in remission [31] and were markedly higher than those of a general

community sample [41, 42]. This suggests that occupational stress and chronic disruption of daily routines may significantly impair rhythm regulation. Notably, the presence of substantial rhythm dysregulation among participants without current depressive episodes points to a latent, subthreshold dysfunction that may represent an early expression of rhythm instability within the DYMERS continuum.

Consistent with this interpretation, our results can be conceptually situated within a stress-rhythm dysregulation-DYMERS-mood disorder pathway model, whereby chronic stress exposure contributes to rhythm desynchronization, which in turn increases vulnerability to mood dysregulation. Future longitudinal work could visually summarize this conceptual pathway to better illustrate the progression from occupational stress to potential clinical outcomes.

#### 4.1. Implications for Research

Our study provides heuristic evidence for rhythm dysregulation and its co-occurrence with depressive symptoms in HCWs, but the cross-sectional design precludes causal inference. Longitudinal and experimental studies are needed to clarify whether rhythm dysregulation represents a true risk factor or an early manifestation of emerging mood pathology. Future research should also explore the scalability, cultural adaptability, and efficacy of rhythm-based interventions, and examine potential moderators such as individual vulnerability, work schedules, and organizational context.

#### 4.2. Implications for Clinical Practice

The clinical implications of our findings are considerable. Research on preventive strategies targeting rhythm dysregulation is still limited, with only a few small pilot studies available [41, 42]. Our study supports the concept of DYMERS as a prodromal syndrome characterized by rhythm dysregulation, which may represent a relevant clinical target in high-risk populations such as healthcare workers (HCWs). Given the robust evidence for Interpersonal and Social Rhythm Therapy (IPSRT) in stabilizing daily rhythms and reducing relapse in BD [43-45], adapting rhythm-focused psychoeducational interventions to HCWs appears promising. This alignment reinforces the translational potential of the current data, suggesting that early rhythm-stabilizing programs could serve as low-intensity, scalable preventive tools aimed at enhancing resilience and mitigating the progression toward mood disorders. While organizational and structural factors (*e.g.*, adequate staffing, professional recognition, workload management) remain essential determinants of healthcare workers' psychological well-being, individual-level rhythm stabilization strategies may offer additional, complementary benefits. We propose a flexible, structured intervention aimed at promoting rhythm regulation, mood stabilization, and stress management. The program combines psychoeducation, practical skills, and home exercises and can be tailored to individual or group needs. Key elements include the following:

- DYMERS framework and vulnerability factors, including the brief psychoeducation on DYMERS as a heuristic model linking behavioral rhythm dysregulation to mood vulnerability [37, 46, 47], as well as discussion of genetic, environmental, and lifestyle moderators (including heritability of hyperactivity-exploratory traits) and their relevance in healthcare settings [13, 46, 48-51].
- Sleep-wake hygiene and daily routine structuring [52-58] and personal stressors and specific mood triggers [36, 59, 60].
- Stress and emotion regulation and cognitive strategies, including mindfulness and relaxation techniques [33, 61-65].
- Social and interpersonal skill development to maintain stable routines and support networks [43, 63, 66-72].
- Early warning signs monitoring and mood tracking to enhance self-awareness and proactive coping [73-77].

This approach is intended as a conceptual and heuristic framework rather than a validated clinical protocol. It highlights how rhythm-oriented interventions could complement organizational strategies (*e.g.*, staffing, workload management) to reduce HCWs' psychological distress and enhance resilience. It aims to guide both empirical research and clinical translation toward rhythm-oriented prevention and early detection strategies.

### 5. LIMITATIONS

Despite the promising findings, this study has several limitations that should be acknowledged.

First, the voluntary convenience sampling and relatively small sample size may limit the generalizability of the results. The recruitment strategy may have introduced self-selection bias, as individuals experiencing higher occupational stress might have been less likely to participate. Second, the cross-sectional design precludes any conclusions regarding causality or temporal dynamics, underscoring the need for longitudinal follow-up studies to examine potential trajectories toward mood or circadian rhythm disorders. Third, because participants were recruited within a single healthcare organization, it was not possible to collect certain socio-demographic data (such as years of service, department, or shift type) without compromising anonymity. While this choice ensured compliance with privacy and ethical principles, it limits the ability to explore potential moderators or sources of variability in the outcomes. Fourth, the reliance on self-reported measures may have introduced response bias and measurement inaccuracy, particularly concerning subjective experiences of rhythm dysregulation or mental health symptoms.

Finally, this study represents a secondary and exploratory analysis within a broader research framework, primarily intended to generate hypotheses and identify potential risk factors for mental health vulnerability among healthcare workers. Within this heuristic perspective, a preliminary preventive intervention model was proposed not as a validated clinical tool, but as a conceptual framework aimed at guiding future empirical research and translational efforts toward the development

of evidence-based prevention and early intervention strategies for this at-risk population.

## CONCLUSION

While this single-center study conducted within the Italian healthcare context limits external validity, the findings contribute to understanding rhythm dysregulation as a potential early indicator of mental health vulnerability in healthcare workers. Future research should include larger and more heterogeneous samples, adopt longitudinal designs to monitor progression toward mood disorders, and rigorously evaluate the effectiveness of rhythm regulation interventions within preventive psychiatry frameworks. The integration of digital tools for continuous and objective monitoring of social and biological rhythms may further enhance assessment precision and personalization.

From an institutional perspective, these findings also align with Italian occupational health policies, which mandate preventive and educational actions to safeguard workers' psychological well-being. Incorporating rhythm-stabilizing and mental health literacy programs within workplace training could therefore represent a feasible and legally supported avenue for early prevention in healthcare settings [78].

## AUTHORS' CONTRIBUTIONS

The contributions of the authors are as follows: E.C., A.U., C.A.G., and G.C.: Conceptualized and designed the study; E.C., S.L., M.A., and S.S.: Completed the investigation; G.C.: was responsible for data collection and management; C.A.G., R.R.R., A.R.R., and M.G.C.: Contributed to data analysis and interpretation; E.C. and A.P.: Drafted the manuscript. All authors critically revised the manuscript for intellectual content and approved the final version. A.P.: Supervised the project and served as the corresponding author.

## LIST OF ABBREVIATIONS

HCWs	= Healthcare Workers
DYMERS	= Dysregulation of Mood, Energy, and Social Rhythms
PHQ-9	= Patient Health Questionnaire-9
BRAIN	= Biological Rhythms Interview of Assessment in Neuropsychiatry
IPSRT	= Interpersonal and Social Rhythm Therapy
PTSD	= Post-Traumatic Stress Disorder
ADHD	= Attention Deficit/ Hyperactivity Disorder

## ETHICS APPROVAL AND CONSENT TO PARTICIPATE

The study received approval from the Independent Ethics Committee of the Azienda Mista Ospedaliero Universitaria, Italy (Protocol No. PG/2018/8822), including the subsequent amendment dated 28 May 2023.

## HUMAN AND ANIMAL RIGHTS

Procedures conformed to the principles outlined in the Declaration of Helsinki, including its most recent revision by the World Medical Association (2020).

## CONSENT FOR PUBLICATION

Participants provided written informed consent after being fully informed about the study aims, data confidentiality, and their right to withdraw at any time without penalty. To ensure participant support, a dedicated telephone line and email address were established to answer any questions or concerns related to study participation.

## STANDARDS OF REPORTING

STROBE guidelines were followed.

## AVAILABILITY OF DATA AND MATERIALS

All data generated or analyzed during this study are included in this published article.

## FUNDING

None.

## CONFLICT OF INTEREST

The author, Dr. Mauro Giovanni Carta, is the EIC of the journal *Clinical Practice & Epidemiology in Mental Health*.

## ACKNOWLEDGEMENTS

Declared none.

## REFERENCES

- [1] Ehlers CL, Frank E, Kupfer DJ. Social zeitgebers and biological rhythms. A unified approach to understanding the etiology of depression. *Arch Gen Psychiatry* 1988; 45(10): 948-52. <http://dx.doi.org/10.1001/archpsyc.1988.01800340076012> PMID: 3048226
- [2] Kahawage P, Bullock B, Meyer D, *et al.* Social rhythm disruption is associated with greater depressive symptoms in people with mood disorders: Findings from a multinational online survey during COVID-19. *Can J Psychiatry* 2022; 67(11): 832-41. <http://dx.doi.org/10.1177/07067437221097905> PMID: 35535550
- [3] Bedrosian TA, Nelson RJ. Timing of light exposure affects mood and brain circuits. *Transl Psychiatry* 2017; 7(1): e1017-7. <http://dx.doi.org/10.1038/tp.2016.262> PMID: 28140399
- [4] Alloy LB, Abramson LY, Urosevic S, Walshaw PD, Nusslock R, Neeren AM. The psychosocial context of bipolar disorder: Environmental, cognitive, and developmental risk factors. *Clin Psychol Rev* 2005; 25(8): 1043-75. <http://dx.doi.org/10.1016/j.cpr.2005.06.006> PMID: 16140445
- [5] Sperry SH, Boland EM, Burgess HJ, *et al.* Bipolar disorder is characterized by chronotype instability: A longitudinal investigation of circadian typology and mood. *Psychiatry Res* 2024; 340: 116123. <http://dx.doi.org/10.1016/j.psychres.2024.116123> PMID: 39146617
- [6] Gold A, Sylvia L. The role of sleep in bipolar disorder. *Nat Sci Sleep* 2016; 8: 207-14. <http://dx.doi.org/10.2147/NSS.S85754> PMID: 27418862
- [7] Zangani C, Casetta C, Saunders AS, Donati F, Maggioni E, D'Agostino A. Sleep abnormalities across different clinical stages of bipolar disorder: A review of EEG studies. *Neurosci Biobehav Rev* 2020; 118: 247-57.

- <http://dx.doi.org/10.1016/j.neubiorev.2020.07.031> PMID: 32738263
- [8] Plante DT, Winkelman JW. Sleep disturbance in bipolar disorder: Therapeutic implications. *Am J Psychiatry* 2008; 165(7): 830-43. <http://dx.doi.org/10.1176/appi.ajp.2008.08010077> PMID: 18483132
- [9] Crowe M, Beaglehole B, Inder M. Social rhythm interventions for bipolar disorder: A systematic review and rationale for practice. *J Psychiatr Ment Health Nurs* 2016; 23(1): 3-11. <http://dx.doi.org/10.1111/jpm.12271> PMID: 26459928
- [10] Miziou S, Tsitsipa E, Moysidou S, *et al.* Psychosocial treatment and interventions for bipolar disorder: a systematic review. *Ann Gen Psychiatry* 2015; 14(1): 19. <http://dx.doi.org/10.1186/s12991-015-0057-z> PMID: 26155299
- [11] Giovanni Carta M, Kalcev G, Scano A, *et al.* The impact of MDQ positivity on quality of life impairment: Does it support the hypothesis of "Dysregulation of mood, energy, and social rhythms syndrome" (DYMERS)? *J Public Health Res* 2023; 12(4): 22799036231208356. <http://dx.doi.org/10.1177/22799036231208356> PMID: 37927350
- [12] Carta MG, Fornaro M, Primavera D, Nardi AE, Karam E. Dysregulation of mood, energy, and social rhythms syndrome (DYMERS): A working hypothesis. *J Public Health Res* 2024; 13(2): 22799036241248022. <http://dx.doi.org/10.1177/22799036241248022> PMID: 38680762
- [13] Carta MG, Kalcev G, Scano A, *et al.* Is bipolar disorder the consequence of a genetic weakness or not having correctly used a potential adaptive condition? *Brain Sci* 2022; 13(1): 16. <http://dx.doi.org/10.3390/brainsci13010016> PMID: 36671999
- [14] Primavera D, Cossu G, Marchegiani S, Preti A, Nardi AE. Does the dysregulation of social rhythms syndrome (DYMERS) be considered an essential component of panic disorders? *Clin Pract Epidemiol Ment Health* 2024; 20(1): e17450179293272. <http://dx.doi.org/10.2174/0117450179293272240328053722> PMID: 38774791
- [15] Primavera D, Aviles Gonzalez CI, Romano F, *et al.* Does the response to a stressful condition in older adults with life rhythm dysregulations provide evidence of the existence of the "Dysregulation of mood, energy, and social rhythms syndrome"? *Healthcare* 2023; 12(1): 87. <http://dx.doi.org/10.3390/healthcare12010087> PMID: 38200993
- [16] Nash C. An initial scoping review of dysregulation of mood, energy, and social rhythms syndrome (DYMERS) regarding burnout in healthcare professionals during COVID-19. *J Clin Med* 2025; 14(3): 1035. <http://dx.doi.org/10.3390/jcm14031035> PMID: 39941704
- [17] Primavera D, Fornaro M, Carrà G, *et al.* Mood disorder questionnaire positivity in systemic lupus erythematosus and other chronic diseases including screen bipolar disorders or rhythm and energy dysregulation syndromes (DYMERS). *Clin Pract Epidemiol Ment Health* 2024; 20(1): e17450179303653. <http://dx.doi.org/10.2174/0117450179303653240705051227> PMID: 39135945
- [18] Zhao Y, Richardson A, Poyser C, Butterworth P, Strazdins L, Leach LS. Shift work and mental health: a systematic review and meta-analysis. *Int Arch Occup Environ Health* 2019; 92(6): 763-93. <http://dx.doi.org/10.1007/s00420-019-01434-3> PMID: 31055776
- [19] Wu QJ, Sun H, Wen ZY, *et al.* Shift work and health outcomes: An umbrella review of systematic reviews and meta-analyses of epidemiological studies. *J Clin Sleep Med* 2022; 18(2): 653-62. <http://dx.doi.org/10.5664/jcsm.9642> PMID: 34473048
- [20] Brown JP, Martin D, Nagaria Z, Verceles AC, Jobe SL, Wickwire EM. Mental health consequences of shift work: An updated review. *Curr Psychiatry Rep* 2020; 22(2): 7. <http://dx.doi.org/10.1007/s11920-020-1131-z> PMID: 31955278
- [21] Bernstrom VH, Alves DE, Ellingsen D, Ingelsrud MH. Healthy working time arrangements for healthcare personnel and patients: a systematic literature review. *BMC Health Serv Res* 2019; 19(1): 193. <http://dx.doi.org/10.1186/s12913-019-3993-5> PMID: 30917819
- [22] Coelho J, Lucas G, Micoulaud-Franchi JA, *et al.* Sleep timing, workplace well-being and mental health in healthcare workers. *Sleep Med* 2023; 111: 123-32. <http://dx.doi.org/10.1016/j.sleep.2023.09.013> PMID: 37769583
- [23] Liu H, Tao TJ, Chan SKY, *et al.* Daily routine disruptions and psychiatric symptoms amid COVID-19: a systematic review and meta-analysis of data from 0.9 million individuals in 32 countries. *BMC Med* 2024; 22(1): 49. <http://dx.doi.org/10.1186/s12916-024-03253-x> PMID: 38302921
- [24] von Elm E, Altman DG, Egger M, Pocock SJ, Gøtzsche PC, Vandenbroucke JP. The strengthening the reporting of observational studies in epidemiology (STROBE) statement: Guidelines for reporting observational studies. *PLoS Med* 2007; 4(10): e296. <http://dx.doi.org/10.1371/journal.pmed.0040296> PMID: 17941714
- [25] Mazzotti E, Fassone G, Picardi A, Sagoni E, Ramieri L, Lega I. The patient health questionnaire (PHQ) for the screening of psychiatric disorders: A validation study against the Structured Clinical Interview for DSM-IV axis I (SCID-I). *Ital J Psychopathol* 2003; 9(3): 235-42.
- [26] Kroenke K, Spitzer RL, Williams JBW. The PHQ-9. *J Gen Intern Med* 2001; 16(9): 606-13. <http://dx.doi.org/10.1046/j.1525-1497.2001.016009606.x> PMID: 11556941
- [27] Moro MF, Carta MG, Pintus M, *et al.* Validation of the Italian version of the biological rhythms interview of assessment in neuropsychiatry (BRIAN): Some considerations on its screening usefulness. *Clin Pract Epidemiol Ment Health* 2014; 10(1): 48-52. <http://dx.doi.org/10.2174/1745017901410010048> PMID: 24987447
- [28] Rosa AR, Comes M, Torrent C, *et al.* Biological rhythm disturbance in remitted bipolar patients. *Int J Bipolar Disord* 2013; 1(1): 6. <http://dx.doi.org/10.1186/2194-7511-1-6> PMID: 25505673
- [29] Allegra OR, Leng X, Vaccarino A, *et al.* Performance of the biological rhythms interview for assessment in neuropsychiatry: An item response theory and actigraphy analysis. *J Affect Disord* 2018; 225: 54-63. <http://dx.doi.org/10.1016/j.jad.2017.07.047> PMID: 28787704
- [30] Cheong HC, Chau SWH, Ng LY, *et al.* Chinese self-report version of biological rhythms interview for assessment in neuropsychiatry (C-BRIAN-SR) - psychometric properties and prospective follow-up in patients with non-seasonal depression. *Chronobiol Int* 2024; 41(7): 1008-20. <http://dx.doi.org/10.1080/07420528.2024.2373215> PMID: 38953315
- [31] Sancassiani F, Lorrain S, Cossu G, *et al.* The effects of "VelaMente?!" project on social functioning of people with severe psychosocial disabilities. *Clin Pract Epidemiol Ment Health* 2017; 13(1): 220-32. <http://dx.doi.org/10.2174/1745017901713010220> PMID: 29299045
- [32] Carta MG, Ouali U, Perra A, *et al.* Living With Bipolar Disorder in the Time of Covid-19: Biorhythms During the Severe Lockdown in Cagliari, Italy, and the Moderate Lockdown in Tunis, Tunisia. *Front Psychiatry* 2021; 12: 634765. <http://dx.doi.org/10.3389/fpsy.2021.634765> PMID: 33716829
- [33] Sancassiani F, Perra A, Kurtschka PK, *et al.* Improving social and personal rhythm dysregulation in young and old adults with bipolar disorder: Post-hoc analysis of a feasibility randomized controlled trial using virtual reality-based intervention. *J Clin Med* 2024; 13(13): 3786. <http://dx.doi.org/10.3390/jcm13133786> PMID: 38999351
- [34] Chupanich P, Aotprapai P, Seesophon S, Laoraksawong P. Factors associated with stress among healthcare personnel after COVID-19 in Northeast Thailand: A cross-sectional study. *Clin Pract Epidemiol Ment Health* 2024; 20(1): e17450179327231. <http://dx.doi.org/10.2174/0117450179327231240924054645> PMID: 39850106

- [35] Liu Y, Li L, Jiang X, *et al.* Mental state, biological rhythm and social support among healthcare workers during the early stages of the COVID-19 epidemic in Wuhan. *Heliyon* 2022; 8(7): e09439. <http://dx.doi.org/10.1016/j.heliyon.2022.e09439> PMID: 35601229
- [36] Carta MG, Preti A, Akiskal HS. Coping with the new era: Noise and light pollution, hyperactivity and steroid hormones. Towards an evolutionary view of bipolar disorders. *Clin Pract Epidemiol Ment Health* 2018; 14(1): 33-6. <http://dx.doi.org/10.2174/1745017901814010033> PMID: 29541149
- [37] Carta MG, Kalcev G, Fornaro M, *et al.* Does screening for bipolar disorders identify a "Dysregulation of mood, energy, and social rhythms syndrome" (DYMERS)? A heuristic working hypothesis. *J Clin Med* 2023; 12(15): 5162. <http://dx.doi.org/10.3390/jcm12155162> PMID: 37568562
- [38] Asaoka H, Watanabe K, Miyamoto Y, *et al.* Association of depressive symptoms with incidence and mortality rates of COVID-19 over 2 years among healthcare workers in 20 countries: multi-country serial cross-sectional study. *BMC Med* 2024; 22(1): 386. <http://dx.doi.org/10.1186/s12916-024-03585-8> PMID: 39267052
- [39] Rivera-Segarra E, Mascayano F, Alnasser L, *et al.* Global mental health research and practice: A decolonial approach. *Lancet Psychiatry* 2022; 9(7): 595-600. [http://dx.doi.org/10.1016/S2215-0366\(22\)00043-8](http://dx.doi.org/10.1016/S2215-0366(22)00043-8) PMID: 35397800
- [40] Czepiel D, Hoek HW, van der Markt A, *et al.* The association between exposure to COVID-19 and mental health outcomes among healthcare workers. *Front Public Health* 2022; 10: 896843. <http://dx.doi.org/10.3389/fpubh.2022.896843> PMID: 35757645
- [41] Goldstein TR, Fersch-Podrat R, Axelson DA, *et al.* Early intervention for adolescents at high risk for the development of bipolar disorder: Pilot study of Interpersonal and Social Rhythm Therapy (IPSRT). *Psychotherapy (Chic)* 2014; 51(1): 180-9. <http://dx.doi.org/10.1037/a0034396> PMID: 24377402
- [42] Goldstein TR, Merranko J, Krantz M, *et al.* Early intervention for adolescents at-risk for bipolar disorder: A pilot randomized trial of Interpersonal and Social Rhythm Therapy (IPSRT). *J Affect Disord* 2018; 235: 348-56. <http://dx.doi.org/10.1016/j.jad.2018.04.049> PMID: 29665518
- [43] Frank E, Swartz HA, Boland E. Interpersonal and social rhythm therapy: An intervention addressing rhythm dysregulation in bipolar disorder. *Dialogues Clin Neurosci* 2007; 9(3): 325-32. <http://dx.doi.org/10.31887/DCNS.2007.9.3/efrank> PMID: 17969869
- [44] Colom F. Keeping therapies simple: Psychoeducation in the prevention of relapse in affective disorders. *Br J Psychiatry* 2011; 198(5): 338-40. <http://dx.doi.org/10.1192/bjp.bp.110.090209> PMID: 21525516
- [45] Miklowitz DJ. *Bipolar disorder: A family-focused treatment approach*. New York: The Guilford Press 2008. <https://www.amazon.com/Bipolar-Disorder-Family-Focused-Treatment-Approach/dp/1572302836>
- [46] Craddock N, Sklar P. Genetics of bipolar disorder. *Lancet* 2013; 381(9878): 1654-62. [http://dx.doi.org/10.1016/S0140-6736\(13\)60855-7](http://dx.doi.org/10.1016/S0140-6736(13)60855-7) PMID: 23663951
- [47] Türközer HB, Guvenek-Cokol PE, Millman ZB, Dickstein D, Öngür D. Familial high-risk for psychosis and bipolar disorder amongst youth in an academic healthcare system: A real-world assessment of clinical characteristics and implications for clinical care. *Early Interv Psychiatry* 2025; 19(2): e70019. <http://dx.doi.org/10.1111/eip.70019> PMID: 39925211
- [48] Kalcev G, Scano A, Orrù G, *et al.* Is a genetic variant associated with bipolar disorder frequent in people without bipolar disorder but with characteristics of hyperactivity and novelty seeking? *Clin Pract Epidemiol Ment Health* 2023; 19(1): e174501792303280. <http://dx.doi.org/10.2174/17450179-v19-e230419-2022-53> PMID: 37916199
- [49] Kalcev G, Cossu G, Preti A, *et al.* Development and validation of the questionnaire for adaptive hyperactivity and goal achievement (AHGA). *Clin Pract Epidemiol Ment Health* 2023; 19(1): e174501792303281. <http://dx.doi.org/10.2174/17450179-v19-e230419-2022-50> PMID: 37916197
- [50] Ouali U, Aissa A, Rejaibi S, *et al.* Hyperactivity and risk for dysregulation of mood, energy, and social rhythms syndrome (DYMERS): Standardization of a simple one-item screener versus the mood disorder questionnaire (MDQ). *J Clin Med* 2024; 13(15): 4433. <http://dx.doi.org/10.3390/jcm13154433> PMID: 39124700
- [51] Tyrer P, Yang M, Tyrer H, Malhi G. Factors predisposing to the onset of bipolar disorder: A 30-Year longitudinal study. *Bipolar Disord* 2025; 27(4): 310-5. <http://dx.doi.org/10.1111/bdi.70026> PMID: 40152415
- [52] Haynes PL, Gengler D, Kelly M. Social rhythm therapies for mood disorders: An update. *Curr Psychiatry Rep* 2016; 18(8): 75. <http://dx.doi.org/10.1007/s11920-016-0712-3> PMID: 27338753
- [53] Boland EM, Goldschmied JR, Kelly MR, Perkins S, Gehrman PR, Haynes PL. Social rhythm regularity moderates the relationship between sleep disruption and depressive symptoms in veterans with post-traumatic stress disorder and major depressive disorder. *Chronobiol Int* 2019; 36(10): 1429-38. <http://dx.doi.org/10.1080/07420528.2019.1644344> PMID: 31368369
- [54] De Pasquale C, El Kazzi M, Sutherland K, *et al.* Sleep hygiene: What do we mean? A bibliographic review. *Sleep Med Rev* 2024; 75: 101930. <http://dx.doi.org/10.1016/j.smrv.2024.101930> PMID: 38761649
- [55] Murray G, Gottlieb J, Swartz HA. Maintaining daily routines to stabilize mood: Theory, data, and potential intervention for circadian consequences of COVID-19. *Can J Psychiatry* 2021; 66(1): 9-13. <http://dx.doi.org/10.1177/0706743720957825> PMID: 32909832
- [56] Arlinghaus KR, Johnston CA. The importance of creating habits and routine. *Am J Lifestyle Med* 2019; 13(2): 142-4. <http://dx.doi.org/10.1177/1559827618818044> PMID: 30800018
- [57] Alfian SD, Thurfah JN, Griselda M, Puspitasari IM. Sleep disturbances and depression levels among general Indonesian population: A national survey. *Clin Pract Epidemiol Ment Health* 2024; 11(20): e17450179326359. <http://dx.doi.org/10.2174/0117450179326359240903045716> PMID: 39850107
- [58] Zou H, Zhou H, Yan R, Yao Z, Lu Q. Chronotype, circadian rhythm, and psychiatric disorders: Recent evidence and potential mechanisms. *Front Neurosci* 2022; 16: 811771. <http://dx.doi.org/10.3389/fnins.2022.811771> PMID: 36033630
- [59] Riachi E, Holma J, Laitila A. Psychotherapists' views on triggering factors for psychological disorders. *Discover Psychology* 2022; 2(1): 44. <http://dx.doi.org/10.1007/s44202-022-00058-y> PMID: 40477882
- [60] Kozubal M, Szuster A, Wielgopalan A. Emotional regulation strategies in daily life: the intensity of emotions and regulation choice. *Front Psychol* 2023; 14: 1218694. <http://dx.doi.org/10.3389/fpsyg.2023.1218694> PMID: 37645071
- [61] Saccaro LF, Giff A, De Rossi MM, Piquet C. Interventions targeting emotion regulation: A systematic umbrella review. *J Psychiatr Res* 2024; 174: 263-74. <http://dx.doi.org/10.1016/j.jpsychires.2024.04.025> PMID: 38677089
- [62] Nowakowska-Domagala K, Podlecka M, Sadowski K, Pietras T, Mokros L. The relationship between chronotype, dispositional mindfulness and suicidal ideation among medical students: Mediating role of anxiety, insomnia and social dysfunction. *J Sleep Res* 2023; 32(4): e13823. <http://dx.doi.org/10.1111/jsr.13823> PMID: 36682738
- [63] Alloy LB, Walsh RFL, Smith LT, *et al.* Circadian, reward, and emotion systems in teens prospective longitudinal study: Protocol overview of an integrative reward-circadian rhythm model of first onset of bipolar spectrum disorder in adolescence. *BMC*

- Psychiatry 2023; 23(1): 602.  
<http://dx.doi.org/10.1186/s12888-023-05094-z> PMID: 37592214
- [64] Guendelman S, Medeiros S, Rampes H. Mindfulness and Emotion Regulation: Insights from Neurobiological, Psychological, and Clinical Studies. *Front Psychol* 2017; 8: 220.  
<http://dx.doi.org/10.3389/fpsyg.2017.00220> PMID: 28321194
- [65] Flores-Kanter PE, Moretti L, Medrano LA. A narrative review of emotion regulation process in stress and recovery phases. *Heliyon* 2021; 7(6): e07218.  
<http://dx.doi.org/10.1016/j.heliyon.2021.e07218> PMID: 34179528
- [66] Sylvia LG, Alloy LB, Hafner JA, Gauger MC, Verdon K, Abramson LY. Life events and social rhythms in bipolar spectrum disorders: a prospective study. *Behav Ther* 2009; 40(2): 131-41.  
<http://dx.doi.org/10.1016/j.beth.2008.04.003> PMID: 19433144
- [67] Picardi A, Filastro A. Development and validation of a self-report measure of existential well-being. *Clin Pract Epidemiol Ment Health* 2025; 21(1): e17450179366317.  
<http://dx.doi.org/10.2174/0117450179366317250410071321> PMID: 40688403
- [68] Frank E, Swartz HA, Kupfer DJ. Interpersonal and social rhythm therapy: managing the chaos of bipolar disorder. *Biol Psychiatry* 2000 Sep 15; 48(6): 593-604.  
[http://dx.doi.org/10.1016/s0006-3223\(00\)00969-0](http://dx.doi.org/10.1016/s0006-3223(00)00969-0) PMID: 11018230
- [69] McMurrin M, Crawford MJ, Reilly J, *et al.* Psychoeducation with problem-solving (PEPS) therapy for adults with personality disorder: A pragmatic randomised controlled trial to determine the clinical effectiveness and cost-effectiveness of a manualised intervention to improve social functioning. *Health Technol Assess* 2016; 20(52): 1-250.  
<http://dx.doi.org/10.3310/hta20520> PMID: 27431341
- [70] Smits M, Kim CM, van Goor H, Ludden GDS. From digital health to digital well-being: Systematic scoping review. *J Med Internet Res* 2022; 24(4): e33787.  
<http://dx.doi.org/10.2196/33787> PMID: 35377328
- [71] Rabelo JL, Cruz BF, Ferreira JDR, Viana BM, Barbosa IG. Psychoeducation in bipolar disorder: A systematic review. *World J Psychiatry* 2021; 11(12): 1407-24.  
<http://dx.doi.org/10.5498/wjp.v11.i12.1407> PMID: 35070785
- [72] Seyyedi Nasooh Abad M, Aemmi SZ, Turati G, Esmaily H. Effectiveness of the motivational interviewing technique on self-efficacy and self-esteem in women facing/experiencing intimate partner violence. *Arch Psychiatr Nurs* 2025; 57: 151938.  
<http://dx.doi.org/10.1016/j.apnu.2025.151938> PMID: 40816774
- [73] Maassen EF, Maathuis L, Regeer BJ, Kupka RW, Regeer EJ. Prodromal symptoms of a first manic episode: a qualitative study to the perspectives of patients with bipolar disorder and their caregivers'. *Int J Bipolar Disord* 2024; 12(1): 38.  
<http://dx.doi.org/10.1186/s40345-024-00360-9> PMID: 39542978
- [74] Álvarez-Cadenas L, García-Vázquez P, Ezquerro B, *et al.* Detection of bipolar disorder in the prodromal phase: A systematic review of assessment instruments. *J Affect Disord* 2023; 325: 399-412.  
<http://dx.doi.org/10.1016/j.jad.2023.01.012> PMID: 36623571
- [75] Roncone R, Mazza M, Ussorio D, *et al.* The questionnaire of family functioning: A preliminary validation of a standardized instrument to evaluate psychoeducational family treatments. *Community Ment Health J* 2007; 43(6): 591-607.  
<http://dx.doi.org/10.1007/s10597-007-9093-8> PMID: 17619149
- [76] Falloon IRH, Held T, Roncone R, Coverdale JH, Laidlaw TM. Optimal treatment strategies to enhance recovery from schizophrenia. *Aust N Z J Psychiatry* 1998; 32(1): 43-9.  
<http://dx.doi.org/10.3109/00048679809062704> PMID: 9565182
- [77] Fristad MA, Gavazzi SM, Mackinaw-Koons B. Family psychoeducation. *Biol Psychiatry* 2003; 53(11): 1000-8.  
[http://dx.doi.org/10.1016/S0006-3223\(03\)00186-0](http://dx.doi.org/10.1016/S0006-3223(03)00186-0) PMID: 12788245
- [78] World medical association declaration of Helsinki. *JAMA* 2025; 333(1): 71-4.  
<http://dx.doi.org/10.1001/jama.2024.21972> PMID: 39425955